

PATIENT MEDICATION INFORMATION

READ THIS FOR SAFE AND EFFECTIVE USE OF YOUR MEDICINE

PrCITALOPRAM

Citalopram Tablets

Read this carefully before you start taking **CITALOPRAM** and each time you get a refill. This leaflet is a summary and will not tell you everything about this drug. Talk to your healthcare professional about your medical condition and treatment and ask if there is any new information about **CITALOPRAM**.

What is CITALOPRAM used for?

CITALOPRAM is used to relieve the symptoms of depression in adults. Your doctor will keep evaluating if CITALOPRAM is still safe and effective for you if you take it for a long time.

How does CITALOPRAM work?

CITALOPRAM is known as an antidepressant and belongs to a group of medicines called Selective Serotonin Reuptake Inhibitors (SSRIs).

CITALOPRAM works by increasing the levels of a chemical in the brain called serotonin.

What are the ingredients in CITALOPRAM:

Medicinal ingredient: citalopram hydrobromide

Non-medicinal ingredients: Colloidal silicon dioxide, hydroxypropyl cellulose, hydroxypropyl methylcellulose, lactose monohydrate, magnesium stearate, microcrystalline cellulose, polyethylene glycol, sodium starch glycolate and titanium dioxide.

CITALOPRAM comes in the following dosage forms:

As tablets containing 10 mg, 20 mg or 40 mg citalopram (as citalopram hydrobromide)

Do not use CITALOPRAM if:

- you are allergic to citalopram hydrobromide
- you are allergic to any of the other ingredients in CITALOPRAM or to a component of the container.
- you are also taking the medicine pimozide, used to treat schizophrenia.
- you are currently taking or have recently taken medicines called monoamine oxidase antidepressants such as phenelzine sulphate, tranylcypromine or moclobemide, or other monoamine oxidase inhibitors such as linezolid, methylene blue, selegiline.
- you have been told that you have QT interval prolongation (abnormal electrical activity of the heart) or have been diagnosed with a condition called a congenital long QT syndrome

To help avoid side effects and ensure proper use, talk to your healthcare professional before

you take CITALOPRAM. Talk about any health conditions or problems you may have, including if you:

- have heart problems.
- have diabetes. CITALOPRAM may make it more difficult to control your blood sugar.
- have liver or kidney problems.
- have or have had a seizure disorder.
- have or have had manic episodes or have been diagnosed with bipolar disorder .
- are receiving electroconvulsive therapy.
- have a bleeding disorder or have been told that you have low platelets.
- have a family history of QT/QTc prolongation (abnormal electrical activity of the heart).
- have electrolyte disturbances like low blood potassium, magnesium, or calcium levels or conditions that could lead to this such as vomiting, diarrhea, dehydration.
- had a recent bone fracture or were told you have osteoporosis or risk factors for osteoporosis.
- are taking or have taken any medications (prescription or non-prescription) within the last 14 days, especially monoamine oxidase inhibitors, pimozide, any other antidepressants, triptans used to treat migraines, lithium, opioids (including to treat pain, or drug dependence), or drugs containing tryptophan.
- have habits of alcohol and /or street drug consumption.
- are taking St. John's Wort, an herbal product used to treat depression.

Other warnings you should know about:

- It is important that you and your doctor talk regularly about how you are feeling while you are taking CITALOPRAM.
- CITALOPRAM should not be used in children and adolescents under 18 years of age.

New or Worsened Emotional or Behavioural Problems

When you first start taking CITALOPRAM, or when you dose is changed, you might feel worse instead of better. You may get new or worsened feelings of agitation, anxiety, hostility or impulsivity.

Self-harm and Suicide

Suicidal thoughts and actions can occur in any age group but may be more likely in patients 18 to 24 years old. If you have thoughts of harming or killing yourself at any time, contact your doctor or go to a hospital **right away**. This is more likely if you have had thoughts of harming or killing yourself in the past. Tell your doctor if you have had these thoughts before. This way, they will monitor you more closely while you are taking CITALOPRAM.

You may also find it helpful to tell a relative or close friend that you are depressed. Ask them to read this leaflet. Ask them to tell you if they think your depression is getting worse, or if they

are worried about changes in your behaviour. Seek medical help if they notice these getting

worse.

Activation of Mania

Tell your doctor if you have or have had manic episodes in the past or if you have been diagnosed with bipolar disorder. CITALOPRAM should be used with caution if you have a history of manic episodes. Some patients with bipolar disorder (also known as manic depression) may enter into a manic phase when they start taking CITALOPRAM. Tell your doctor if you experience symptoms of mania such as excessive physical activity, overactive behaviour or thoughts, increased energy, trouble sleeping, racing thoughts, reckless behaviour, excessive happiness or irritability, talking more or faster than usual.

Bleeding Problems

Before taking CITALOPRAM tell your doctor if you have a bleeding disorder including low blood platelets. Drugs from the class that CITALOPRAM belongs to may increase the chance of a bleeding event such as nose bleeds, bruising and even life threatening bleeding. This is more likely if you have a history of a bleeding disorder or are taking other drugs that are known to affect blood clotting. Talk to your doctor about drugs that might increase bleeding.

Pregnancy

Before taking CITALOPRAM, tell your doctor if you are pregnant, think you might be pregnant or are planning to become pregnant. You should not take CITALOPRAM if you are pregnant unless you and your doctor have discussed the risks and decided that you should. Tell your doctor right away if you become pregnant while taking CITALOPRAM. If you take CITALOPRAM near the end of your pregnancy, you could have heavy vaginal bleeding shortly after giving birth.

Effects on Newborns

Some newborn babies whose mothers took medications such as CITALOPRAM during pregnancy have developed problems at birth. These problems include prolonged hospitalisation, breathing support and tube feeding. Symptoms can include:

- feeding and/or breathing difficulties
- bluish skin
- seizures
- body temperature changes
- vomiting
- low blood sugar
- tense or overly relaxed muscles
- vivid reflexes
- tremor
- jitteriness
- irritability
- sleeping difficulties and constant crying.

In most cases, these medications were taken during the third trimester of pregnancy. These

symptoms are caused by the medication itself or from the effects of suddenly stopping the medication. These symptoms normally go away over time. However, if your baby experiences any of these symptoms, contact your doctor as soon as you can.

Persistent Pulmonary Hypertension of the Newborn (PPHN)

If you take CITALOPRAM towards the end of your pregnancy, your newborn may be at risk of having a serious lung condition called Persistent Pulmonary Hypertension of the Newborn (PPHN). This causes breathing problems in newborns soon after birth. Newborn babies may breathe faster and appear bluish. These symptoms usually begin during the first 24 hours after the baby is born. If this happens to your newborn baby, get immediate medical help for them.

Breastfeeding

Tell your doctor if you are breastfeeding or planning to breastfeed. CITALOPRAM is released into breast milk. It is not known if this is safe for your baby. You should not breastfeed a baby if you are taking CITALOPRAM unless you and your doctor have discussed the risks and decided that you should.

Effect on the electrical activity of the heart

CITALOPRAM has an effect on the electrical activity of the heart known as QT/QTc prolongation (abnormal electrical activity of the heart). This can lead to disturbances in heart rhythm (arrhythmias/dysrhythmias) that could result in dizziness, palpitations (sensation of rapid, pounding, or irregular heartbeat), fainting or cardiac arrest and death. This is more likely in patients with risk factors, such as heart disease, heart attack, or in the presence of certain drugs that could interact with the activity of the heart. If you experience any symptoms of a possible heart rhythm disturbance (abnormal heart rate or rhythm), such as dizziness, palpitations (sensation of rapid, pounding, or irregular heartbeat), fainting, or seizures, you should seek immediate medical attention.

Serotonin Toxicity

Serotonin toxicity (also known as Serotonin syndrome): CITALOPRAM can cause Serotonin toxicity, a rare but potentially life-threatening condition. It can cause serious changes in how your brain, muscles and digestive system work. You may develop Serotonin toxicity if you take CITALOPRAM with certain anti-depressants or migraine medications.

Serotonin toxicity symptoms include:

- fever, sweating, shivering, diarrhea, nausea, vomiting;
- muscle shakes, jerks, twitches or stiffness, overactive reflexes, loss of coordination
- fast heartbeat, changes in blood pressure;
- confusion, agitation, restlessness, hallucinations, mood changes, unconsciousness, and coma

Effects on Sexual Function

Taking medicines like CITALOPRAM may cause symptoms of sexual dysfunction. In some cases these symptoms have continued after stopping CITALOPRAM treatment. Talk to your healthcare professional if you experience symptoms such as a decrease in sexual desire, performance or satisfaction.

Risk of Bone Fractures

Taking CITALOPRAM may increase your risk of breaking a bone if you are elderly, have osteoporosis or other risk factors for breaking a bone. You should take extra care to avoid falls especially if you get dizzy or have low blood pressure.

Angle-closure Glaucoma

CITALOPRAM can cause dilation of the pupil. This may cause an acute glaucoma attack in an individual with narrow ocular angles. Having your eyes examined before you take CITALOPRAM could help identify if you are at risk of having angle-closure glaucoma. Get immediate medical help if you experience:

- eye pain
- changes in vision
- swelling or redness in or around the eye.

Driving and using machines

CITALOPRAM may impair your ability to drive or to use machines. Wait until you know how CITALOPRAM affects you before driving or using machines. Do not drive or use machines if CITALOPRAM impairs your ability to do so safely.

Discontinuation Symptoms

Contact your doctor before stopping or reducing your dosage of CITALOPRAM. If you stop or reduce the dosage of CITALOPRAM abruptly, or if you miss a dose, you may experience symptoms such as dizziness, sleep disturbances, abnormal dreams, sensory disturbance like electric shock sensations, agitation, anxiety, headache, tremor (shakiness), nausea, vomiting. Tell your doctor immediately if you have these or any other symptoms. Your doctor may adjust the dosage of CITALOPRAM to reduce the symptoms.

Tell your healthcare professional about all the medicines you take, including any drugs, vitamins, minerals, natural supplements or alternative medicines.

Serious Drug Interactions

Do not use CITALOPRAM if you are taking or have recently taken:

- A monoamine oxidase inhibitor such as phenelzine, tranylcypromine, moclobemide or selegiline, linezolid (as antibiotic) or Methylene blue (intravenous)
- Pimozide

The following may interact with CITALOPRAM:

- drugs to treat heart rhythm disturbances (antiarrhythmics)
- antipsychotics, used to treat schizophrenia
- opioids (including for pain, drug dependence or anesthesia) such as methadone, buprenorphine, tramadol, fentanyl, tapentadol, meperidine or pentazocine
- drugs to treat infections
- drugs to treat nausea and vomiting
- cancer drugs
- asthma drugs
- diuretics (water pills)
- carbamazepine used to treat seizures
- Other SSRIs e.g., escitalopram or any other antidepressant (e.g., imipramine, desipramine)
- lithium, used to treat mood disorder
- tryptophan, for sleep aid or treating anxiety
- cimetidine, for acidity problems
- triptans (e.g., sumatriptan, zolmitriptan, naratriptan), for migraine
- fluconazole, ketoconazole, itraconazole, for treating fungal infection
- erythromycin, used to treat infections
- warfarin, used to prevent clot of blood
- omeprazole, used to treat stomach problems
- any herbal product such as St. John's Wort
- certain medicines which may affect blood clotting and increase bleeding, such as oral anticoagulants (e.g., warfarin, dabigatran), acetylsalicylic acid and other non-steroidal anti-inflammatory drugs (ibuprofen).
- certain medicines used to treat cough, such as dextromethorphan.

Avoid drinking alcohol while taking CITALOPRAM

How to take CITALOPRAM:

- Take CITALOPRAM exactly as your doctor has told you.
- Swallow tablets whole with water. Do not chew them.
- You can take CITALOPRAM with or without food.
- Take CITALOPRAM once a day at the same time every day.
- Continue taking CITALOPRAM even if you do not feel better. It may take several weeks for it to work and improvement may be gradual.
- Keep taking CITALOPRAM for as long as your doctor recommends. Do not stop taking CITALOPRAM abruptly even if you feel better, unless your doctor has told you to.
- Never take more CITALOPRAM than your doctor has prescribed for you.
- Follow all instructions given to you by your doctor.

Usual dose:

The usual dose is one 20 mg tablet once a day. Your doctor might prescribe a lower dose if you are elderly, have liver problems or in other situations.

Overdose:

If you think you, or a person you are caring for, have taken too much CITALOPRAM, contact a healthcare professional, hospital emergency department, or regional poison control centre immediately, even if there are no symptoms.

Missed Dose:

If you forget a dose, take the next dose as planned. Do not take a double dose to make up for a forgotten dose.

What are possible side effects from using CITALOPRAM?

These are not all the possible side effects you may have when taking CITALOPRAM. If you experience any side effects not listed here, tell your healthcare professional.

Side effects may include:

- Fatigue
- Dry mouth
- Increased sweating
- Tremor (shakiness)
- Nausea
- Diarrhea
- Somnolence (sleepiness)
- Ejaculation disorder

Serious side effects and what to do about them			
Symptom / effect	Talk to your healthcare professional		Stop taking drug and get immediate medical help
	Only if severe	In all cases	
UNCOMMON			
Bleeding problems: Bruising or bleeding from the skin, nose or other areas for longer than usual.		✓	
Mania: Excessive physical activity, overactive behaviour or thoughts, increased energy, trouble sleeping, racing thoughts, reckless behaviour, excessive happiness or irritability, talking more or faster than usual.		✓	

Serious side effects and what to do about them			
Symptom / effect	Talk to your healthcare professional		Stop taking drug and get immediate medical help
	Only if severe	In all cases	
RARE			
Gastrointestinal bleeding: Vomiting blood, passing blood in stools		✓	
Angle-closure Glaucoma: (Increased pressure in eyes, change in vision such as hazy or blurred vision) : Eye pain, change in vision, swelling or redness in or around the eye			✓
Low sodium level in blood: Tiredness, weakness, confusion combined with achy, stiff or uncoordinated muscles		✓	
Serotonin Toxicity and Neuroleptic Malignant Syndrome (various symptoms due to high level of serotonin level in the body): A combination of most or all of the following: confusion, restlessness, sweating, shaking, shivering, high fever, hallucinations, sudden jerking of the muscles, muscle stiffness, feeling very agitated or irritable, fast heartbeat. The severity can increase, leading to loss of consciousness.			✓
Hypoglycemia (Low sugar level in blood): Feeling shaky, sweating, chills and clamminess, irritability or impatience, confusion, fast heartbeat, feeling lightheaded or dizzy, nausea, seizure.		✓	
VERY RARE			
Liver disorder: Nausea, vomiting, loss of appetite, itching, yellowing of the skin or eyes, dark urine		✓	
Seizures (fits): Loss of consciousness with uncontrollable shaking			✓
New or Worsened Emotional or Behavioural Problems: Anxiety, hostility or impulsivity Akathisia: Feeling restless and unable to sit or stand still		✓	
Self-harm and suicide: Have thoughts of harming or killing yourself			✓
UNKNOWN			
Heart rhythm disturbance (abnormal heart rate or rhythm): Dizziness, palpitations (sensation of rapid, pounding or irregular heartbeat), fainting, seizures			✓
Postpartum haemorrhage (Heavy vaginal bleeding shortly after birth): Excessive vaginal bleeding after childbirth		✓	

Serious side effects and what to do about them			
Symptom / effect	Talk to your healthcare professional		Stop taking drug and get immediate medical help
	Only if severe	In all cases	
Serious skin reactions: Skin rash, redness of the skin, blistering of the lips, eyes or mouth, skin peeling, fever, chills, headache, cough, body aches			✓
Symptoms after discontinuation or dose reduction: Dizziness, sleep disturbances, abnormal dreams, sensory disturbance like electric shock sensations, agitation, anxiety, headache, tremor (shakiness), nausea, vomiting		✓	

If you have a troublesome symptom or side effect that is not listed here or becomes bad enough to interfere with your daily activities, tell your healthcare professional.

Reporting Side Effects

You can report any suspected side effects associated with the use of health products to Health Canada by:

- Visiting the Web page on Adverse Reaction Reporting (<https://www.canada.ca/en/health-canada/services/drugs-health-products/medeffect-canada/adverse-reaction-reporting.html>) for information on how to report online, by mail or by fax; or
- Calling toll-free at 1-866-234-2345.

NOTE: Contact your health professional if you need information about how to manage your side effects. The Canada Vigilance Program does not provide medical advice.

Storage:

Keep CITALOPRAM out of the reach and sight of children. Store CITALOPRAM at room temperature (15°C - 30°C), in a dry place.

If you want more information about CITALOPRAM:

- Talk to your healthcare professional.
- Find the full product monograph that is prepared for healthcare professionals and includes this Patient Medication Information by visiting the Health Canada website (<https://www.canada.ca/en/health-canada/services/drugs-health-products/drug-products/drug-product-database.html>), or by contacting the sponsor Sivem Pharmaceuticals ULC at: -1-855-788-3153.

This leaflet was prepared by Sivem Pharmaceuticals ULC

Last revised: October 18, 2023